## DENNISON, SCHEINER, SCHULTZ & WAKEMAN 1745 JEFFERSON DAVIS HIGHWAY ARLINGTON, VIRGINIA 22202-3417 2 CRYSTAL SQUARE 4

<u>REMARKS</u>

The claims have been amended to delete all multiple dependencies, and to generally place the claims in better form for US practice.

Respectfully submitted,

Schultz

Registration No. 28666

LAW OFFICES

## APPENDIX

IN THE CLAIMS:

Page 9, lines 1-3: [Patent Claims Method for Adjusting or Controlling a Person's Nutrition and/or Consumption] WHAT IS CLAIMED IS:

- 3. (Amended) Method according to claim 1 [or 2], characterized by the fact that for determining the performance capacity the individual anaerobic threshold of the person is measured or determined.
- 4. (Amended) Method according to [at least one of the previous claims] claim 1, characterized by the fact that for determining the performance capacity a scaling of the performance measured above the individual anaerobic threshold occurs according to the lactate accumulation rate  $\Delta A$ .
- 5. (Amended) Method according to [at least one of the previous claims] claim 1, characterized by the fact that the stress is used as a basis for the IAT and the lactate accumulation rate  $\Delta A$  in determining the nutrition and/or the consumption of a person with regard to his/her carbohydrate and/or fat and/or protein percentages.
- Method according to [at least one of the previous claims] claim 1, characterized by the fact that the individual anaerobic threshold according to Stegmann is used a basis for determining the nutrition and/or consumption of the person

with regard to his/her carbohydrate and/or fat and/or protein percentages.

- 7. (Amended) Method according to [at least one of the previous claims] claim 1, characterized by the fact that when stress occurs in a person over an extended period of time below his/her individual anaerobic threshold, the fat percentage of the nutrition is adjusted comparatively higher than the carbohydrate and the protein percentages.
- (Amended) Method according to [at least one of the previous claims] claim 1, characterized by the fact that with a lactate accumulation rate  $\Delta A$  against  $\Delta A_{max}$  the protein percentage of the nutrition is adjusted up to several times as high as with  $\Delta A = 0$ .
- 9. (Amended) Method according to [at least one of the previous claims] claim 1 for determining the lactate accumulation rate  $\Delta A$ , [characterized by the following procedural steps] comprising the steps of:

measuring the time-dependent lactate concentration change beyond the individual anaerobic threshold,

adjusting a measurement curve to measurement values gained this way, in which the lactate concentration in relation to time is plotted,

determining a first gradient in the measurement curve at

a time  $t_{\text{IAT}}$  that corresponds to the individual anaerobic

703 412-1155

1745 LEFFERSON DAVIS HIGHWAY ARLINGTON, VIRGINIA 22202-3417

threshold,

determining at least one additional gradient in the measurement curve at a time  $t_{x}$  with  $t_{x}$  >  $t_{\text{IAT}}$ 

subtracting the second gradient from the first gradient to determine a difference, which represents the lactate accumulation rate  $\Delta A$ .

10. (Amended) Method according to [at least one of the previous claims] claim 1, characterized by the fact that for determining the performance capacity, different types of stress such as running tests, swimming tests, stepping tests, ergometry methods with graduated or continuous stress increase with and without breaks are used.

Patent Claims

Method for Adjusting or Controlling a Person's Nutrition and/or Consumption

Method for adjusting or controlling the nutrition
 and/or consumption of carbohydrates and/or fats and/or proteins
 in a person subjected to stress,

characterized by the fact

that for the control and/or adjustment of the nutrition and/or consumption of nutrients in a human being his/her performance capacity is determined by determining characteristic performance capacity parameters and that as a function upon the determined performance capacity the carbohydrate and/or fat and/or protein percentage requirements are determined and/or their consumption by a person, while basing the calculation on the stress, which is decisive for the nutrition and/or consumption.

2. Method according to claim 1, characterized by the fact that, for determining the performance capacity, the heart rate and/or blood pressure and/or ergospirometric parameters and/or lactate concentration in the blood are measured or determined as a function upon the stress.

10015445 1515 1015

5

703 412-1155

DENNISON, SCHEINER, SCHULTZ & WAKEMAN
612 CRYSTAL SOUARE 4
1745 JEFFERSON DAVIS HIGHWAN
ARLINGTON, VIRGINIA 22202-34

DENNISON,

- 3. Method according to claim 1 or 2, characterized by the fact that for determining the performance capacity the individual anaerobic threshold of the person is measured or determined.
- 4. Method according to at least one of the previous claims, characterized by the fact that for determining the performance capacity a scaling of the performance measured above the individual anaerobic threshold occurs according to the lactate accumulation rate  $\Delta A$ .
- 5. Method according to at least one of the previous claims, characterized by the fact that the stress is used as a basis for the IAT and the lactate accumulation rate  $\Delta A$  in determining the nutrition and/or the consumption of a person with regard to his/her carbohydrate and/or fat and/or protein percentages.
- characterized by the fact

  that the individual anaerobic threshold according to Stegmann is used a basis for determining the nutrition and/or consumption of the person with regard to his/her carbohydrate

Method according to at least one of the previous

and/or fat and/or protein percentages.

- 7. Method according to at least one of the previous claims, characterized by the fact that when stress occurs in a person over an extended period of time below his/her individual anaerobic threshold, the fat percentage of the nutrition is adjusted comparatively higher than the carbohydrate and the protein percentages.
- 9. Method according to at least one of the previous claims for determining the lactate accumulation rate  $\Delta A$ , characterized by the following procedural steps

measuring the time-dependent lactate concentration change beyond the individual anaerobic threshold,

adjusting a measurement curve to measurement values gained this way, in which the lactate concentration in relation to time is plotted,

determining a first gradient in the measurement curve at a

time  $t_{\text{IAT}}$  that corresponds to the individual anaerobic threshold.

determining at least one additional gradient in the measurement curve at a time  $t_{x}$  with  $t_{x}$  >  $t_{\text{IAT}}$ 

subtracting the second gradient from the first gradient to determine a difference, which represents the lactate accumulation rate  $\Delta A$ .

- 10. Method according to at least one of the previous claims, characterized by the fact that for determining the performance capacity, different types of stress such as running tests, swimming tests, stepping tests, ergometry methods with graduated or continuous stress increase with and without breaks are used.
- 11. Method for adjusting and/or controlling the nutrition and/or consumption of carbohydrates and/or fats and/or proteins of a person who is subjected to stress characterized by the fact that for the control and/or adjustment of the nutrition and/or consumption of nutrients in a human being his/her performance capacity is determined by determining characteristic performance capacity parameters and that the need for and/or consumption of carbohydrates and/or fats and/or proteins in the

03 412-1155

food is determined as a function upon the determined performance capacity of the person, while basing the calculation on the stress-specific substrate mixture ratios that are decisive for the nutrition and/or consumption.

5

13